

FICTITIOUS DOMAIN METHODS TO THE DIRECT NUMERICAL SIMULATION OF THE LIFTING OF ELLIPSOIDS IN THE INCOMPRESSIBLE VISCOUS FLUID

Roland Glowinski^a and Tsorng-Whay Pan^a

^aDepartment of Mathematics
University of Houston
Houston, Texas 77204-3476
roland@math.uh.edu, pan@math.uh.edu

In this talk, we discuss the distributed Lagrange multiplier (DLM)/fictitious domain methods for the direct numerical simulation of the interaction between incompressible viscous fluid and the rigid bodies of general shape, which are moved by the hydrodynamical forces and gravity. We will focus on the lifting of the ellipsoids and the migration of the neutrally buoyant particles in a pressure driven Poiseuille flow in a pipe. For the cases where the particles are neutrally buoyant, a methodology which can handle the difficulties from the DLM approach is presented. We conclude this talk by the presentation of the results of various numerical experiments in two and three dimensions.

References

- [1] R. Glowinski, T.-W. Pan, T.I. Hesla, D.D. Joseph and J. Periaux, A fictitious domain approach to the direct numerical simulation of incompressible viscous flow past moving rigid bodies: Application to particulate flow, J. Comput. Phys., 169 (2001), 363-426.
- [2] T.-W. Pan, R. Glowinski, Direct simulation of the motion of neutrally buoyant circular cylinders in plane Poiseuille flow, J. Comput. Phys., 181 (2002), pp. 260-279.